Sanofi Pasteur's Fluzone® Vaccine Package Insert

Teratogenic effects: Pregnancy Category C:
Animal reproduction studies have not been conducted with
Fluzone vaccine. It is also not known whether Fluzone
vaccine can cause fetal harm when administered to a
pregnant woman or can affect reproduction capacity.
Fluzone vaccine should be given to a pregnant woman only
if clearly needed.

Carcinogenesis, Mutagenesis, Impairment of Fertility: Fluzone vaccine has not been evaluated for carcinogenic or mutagenic potential, or for impairment of fertility.

Under Federal Law (NVICP 1986) your Vaccine Provider is Required to:

- Give you vaccine benefit and risk information prior to vaccination.
- (2) Obtain your written consent before administration.
- (3) Keep a permanent record of all vaccinations given, including the manufacturer's name and lot number.
- (4) Document serious health problems, hospitalizations, injuries and deaths that occur after vaccination in your or your child's permanent medical record.
- (5) Report serious health problems, hospitalizations, injuries and deaths that occur after vaccination to the federal Vaccine Adverse Event Reporting System (VAERS).

Public Schools as Mass Vaccination Centers?

- U.S. Department of Health & Human Services Secretary Kathleen Sebelius is currently mobilizing school superintendents to use public schools as mass vaccination centers.
- U.S. Rep. Gene Green (D-TX) sponsored legislation H.R. 2596 to authorize the use of public schools to carry out mass vaccinations.

WHAT CAN YOU DO? DO SOMETHING!

IMMEDIATELY begin to educate your community about the potential of forced mass vaccination programs and use of public schools for mass vaccination sites.

- Print copies and distribute this flyer.
- Give this information to everyone you know and love: Church members, social groups, school teachers, administrators, first responders (EMTs, paramedics, firemen), local police, sheriff, city council members, county commissioners, chamber of commerce and state legislators. Their support is critical to maintain your right to choose.
- · Write letters to the editor in LOCAL community newspapers.
- DEMAND that public schools are not used as mass vaccination sites.
- Get this information out via the web using Facebook, MySpace or Twitter. Feel free to hot-link or post this information anywhere on the web.

What to do if you Plan to Decline

- Ask your State Attorney General, Legislator or Governor whether your state law exemptions from vaccine programs (medical, religious, philosophical) apply during a declared pandemic or health emergency.
- If exemptions are available, contact your health care or school professionals for forms to fill out giving notice that you or your family members are exempt and are declining the vaccines and/or treatments.
- Find out if swine flu or pandemic vaccines will be required for students and staff in order to attend public schools.
- Pursuant to Federal Law (NVICP 1986), demand that your vaccine provider give you a benefit and risk vaccine information statement prior to vaccination. They must obtain your written consent before administration.
- Find out if exposed persons have the right to self-quarantine at home.

Download and Print Flyer:

Educate yourself



Swine Flu Mass Vaccination Programs - 2009/2010

In the case of the 1976 swine flu pandemic, a government-directed mass vaccination program, several dozen vaccine recipients died and hundreds were injured. Those families who were able to sue and won were awarded millions of dollars for the injuries caused by the untested vaccine formulations.

Three years ago legislators strategically passed The Biodefense and Pandemic Vaccine and Drug Development Act of 2006 (also known as BioShield II). This law strips Americans of the right to a jury trial if harmed by an experimental drug or vaccine during a declared pandemic. Keep in mind there is no guaranteed compensation program for those who may be injured or for families whose loved ones die from being vaccinated or treated.

Thimerosal (Mercury) and other Components

Thimerosal is heavily implicated as a major cause of the mass generational epidemic in developmental disorders (including the autism spectrum disorders, ADD, ADHD and OCD), endocrine-system dysfunction, and immune-system-related chronic diseases of the lungs, bowel, pancreas, liver and cardiovascular system.

Studies show a "... 7 to 1 increase in relative risk of injury for children born to mothers who got a thimerosal-preserved flu shot in the first 4 lunar months of their pregnancy over matched mothers who did not."

(Birth Defects and Drugs in Pregnancy, Heinonem, Slone, Shapiro, 1983)

"Significantly increased risks of autism, speech disorders, mental retardation, personality disorders, thinking abnormalities, ataxia and NDs (neuro-developmental disorders) in general,...were associated with TCV (thimerosal) exposure." (Neuroendocrinology Letters Volume 27 No. 4 2006. Mark R. Geier, M.D., Ph.D. and David A. Geier)

Other components: Formaldehyde (a neurotoxin and carcinogen), egg protein, contaminant viruses, antibiotics, and detergent.

The only Vaccine Information Statement (VIS) that is provided to health professionals who in turn must, by law, provide it to all vaccine recipients, is an official VIS prepared exclusively by the Centers for Disease Control (CDC).

The CDC will not recommend thimerosal- free over thimerosal-containing vaccine to pregnant women, children and the elderly despite the avalanche of independent studies indicating the risks. "No preference is indicated for use of vaccine that does

not contain thimerosal (mercury)...for any group recommended for vaccination including pregnant women."

Not disclosed in the VIS is that the CDC recommendations contradict BOTH the warnings on the manufacturer's package insert and the FDA indications for pregnant manufacturer.

www.cdc.gov/flu/professionals/acip/specificpopulations.htm

CDC: "Vaccination of [all] pregnant women is recommended" by the Advisory Committee on Immunization Practices.

www.cdc.gov/vaccines/pubs/vis/default.htm#flu

FDA: "These are [ACIP] recommendations and are **MOT FDA-approved** indications for use of vaccines during pregnancy", David Horowitz, Esq., Assistant Commissioner for Policy FDA/HHS.

Many Studies Show that Vaccines are Not Effective

CHILDREN: "...no evidence that injecting children 6-24 months of age with a flu shot was any more effective than placebo." Reference: "Vaccines for preventing influenza in healthy children.", The Cochrane Database of Systematic Reviews. 2 (2008).

ASTHMA: "The inactivated flu vaccine does not prevent influenza-related hospitalizations in children, especially the ones with asthma... Children who get the flu vaccine are three times more at risk for hospitalization than children who do not get the vaccine." Reference: The American Thoracic Society's 105th International Conference, May 15-20, 2009, San Diego.

HEALTHY ADULTS: "Vaccination of healthy adults only reduced risk of influenza by 6%..." Reference: Vaccines for preventing influenza in healthy adults. The Cochrane Database of Systematic Reviews. 1(2006).

THE ELDERLY: "Flu shots were non-significant for preventing the flu in elderly living in nursing homes. For elderly living in the community, vaccines were not (significantly) effective against influenza, ILI or pneumonia..." Reference: "Vaccines for preventing influenza in the elderly." The Cochrane Database of Systematic Reviews. 3(2006).

PREGNANT WOMEN: FDA and manufacturer's insert indicate that the Flu shot should NOT be given to pregnant women whether it contains Thimerosal (mercury) or not. Without proof of fetal and reproductive safety, the CDC and Advisory Committee on Immunization Practices (ACIP) recommend that all pregnant women get vaccinated against the flu.

Squalene

Depending on the manufacturer, H1N1 may contain a potentially debilitating oil based adjuvant (additive) primarily composed of squalene (MF59, AS03, AS04).

All laboratory rats injected with squalene (oil) adjuvants developed a disease that left them crippled, dragging their paralyzed hindquarters across their cages. Injected squalene can cause severe arthritis (3 on scale of 4) and severe immune responses such as autoimmune arthritis and lupus.

References: (1) Kenney RT, Edleman, R. "Survey of human-use adjuvants." Expert Review of Vaccines. 2 (2003) p171; (2) Matsumoto, Gary. Vaccine A: The Covert Government Experiment That's Killing Our Soldiers and Why Gl's Are Only the First Victims of this Vaccine. (2004) New York: Basic Books. p54.

Symptoms of Vaccine Injury

- Inability to focus at school and at home. Lack of eye contact.
- Depression, moodiness; tendency to overreact emotionally.
- Delayed or unusual speech patterns such as high pitched or flat intonation; lack of slang or "kidspeak."
- Hyper or hypo sensitivity to light, sound, crowds and other external stimulation.
- Difficulty with fine and gross motor skills such as handwriting, athletic coordination and endurance sports such as running and swimming
- Repetitive behaviors and ritualized activities.
- Difficulty in making and keeping friends later in life.
- Asthma and diet issues.
- Conditions often labeled as ADD/ADHD such as inattention, hyperactivity, and impulsivity.
- Significant abnormal bowel function or vasculitis. (1)
- Birth defects including, cleft palate, microcephaly or pyloric stenosis. (2)
- Guillian Barré Syndrome, a neurological disorder. (3)
- (1) Infection and Autoimmunity by Yehuda Shoenfeld, Noel R. Rose
- (2) American Journal of Epidemiology, "Delivering Influenza Vaccine To Pregnant Women", by David M. Ayoub, MD and F. Edward Yazbak, MD
- (S) U.S. Centers for Disease Control

Who have been Designated as Priority Recipients for the First Wave of Experimental Swine-flu Vaccines?

- School children who have never had a flu shot are targeted for four shots in the fall - twice for seasonal flu, twice for pandemic swine flu. (July 15, 2009 news) Multi-dose vials will contain Thimerosal (49.6-wt-% mercury).
- All students and staff associated with schools (K-12th grade) and children (aged 6 months or older) and staff in child care
- \bullet Pregnant women, children 6 months to 4 years of age, new parents and households in contact with children.
- Health care workers and emergency services sector personnel.
- Adults less than 65 years old with medical conditions which

increase the risk of influenza.

Why haven't our top national leaders been included in this list as first priority recipients?